

Total marks — 100
Attempt ALL questions

Candidates should observe that $g \text{ m s}^{-2}$ denotes the magnitude of the acceleration due to gravity. Where appropriate, take its magnitude to be 9.8 m s^{-2} .

1. A bicycle and rider have a total mass of 70 kg. They are travelling at 12 m s^{-1} . The cyclist applies the brakes for 1.5 seconds, resulting in a total resistive force of 180 newtons.


What is the speed of the bicycle after 1.5 seconds?

3

2. In a children's playground game, four light inextensible ropes are attached at one end to a small toy ring.

Four children each take the other end of a rope and pull it taut.

The ring is in equilibrium and the whole system is in a horizontal plane with appropriate axes as shown in the diagram.



The tensions in the four ropes are P , Q , 80 and 64 newtons respectively, and their directions relative to the axes are shown.

Calculate the magnitude of the tensions P and Q .

4

3. A constant force $\mathbf{F} = (2\mathbf{i} + 3\mathbf{j}) \text{ N}$ acts on a particle as it moves in a straight line from point A to point B with position vectors $(-3\mathbf{i} + \mathbf{j})$ metres and $(6\mathbf{i} + 4\mathbf{j})$ metres respectively.

Calculate the work done by the force.

3

[Turn over

4. Find the equation of the tangent to the curve $y = x \ln x$ at the point where $x = e$. 3

5. The tip of a saw oscillates with simple harmonic motion.

- When the tip is 5 mm from its centre of motion it has a velocity of 2 m s^{-1} .
- When it is 7 mm from the centre it has a velocity of 1 m s^{-1} .

Calculate the amplitude of the motion and find the number of oscillations in one second. 5

6. A remote controlled aircraft is flown from point A to point B. It accelerates for 10 seconds at a constant rate from rest to a take-off speed of 15 m s^{-1} .

Once airborne, it accelerates for a further 20 seconds at a slower constant rate to a cruising speed of $u \text{ m s}^{-1}$.

It maintains this speed for 60 seconds until it lands.

The aircraft then decelerates for 10 seconds to a complete stop.

- (a) Sketch a speed-time graph of the journey, clearly showing all the important information. 2

- (b) (i) If the distance travelled from A to B is 1.725 km, calculate the value of u . 2

- (ii) State one assumption you have made about the path of the aircraft during your calculations. 1


7. An object of mass 9 kg starts from rest at an origin and moves in a straight line so that its acceleration in m s^{-2} is given as $a = 4 - \sqrt{t}$, where t is the time in seconds.

Calculate its maximum speed and hence the increase in kinetic energy. 4

8. (a) Show that $\frac{3x^3 + 8x^2 - 11}{(x+1)(x+3)(x-2)}$ can be written as $3 + \frac{2x^2 + 15x + 7}{x^3 + 2x^2 - 5x - 6}$. 3

- (b) Hence express $\frac{3x^3 + 8x^2 - 11}{(x+1)(x+3)(x-2)}$ in partial fractions. 4

9. A velodrome has a circular track of radius 30 metres, banked at an angle of 32° to the horizontal. The coefficient of friction between a bicycle tyre and the track is 0.3.
- (a) Once the cyclist reaches maximum speed without the bicycle slipping, he cycles for 5 minutes. Assuming he maintains this speed, how many full laps does he complete? 6
- (b) Given that air resistance can be ignored and the cyclist is treated as a particle, what other assumption has been made? 1
10. A stone is thrown from the top of a cliff and the subsequent motion can be modelled in the xy plane by the equations $x=4t$ and $y=20+2t-5t^2$.
- (a) Use parametric differentiation to find $\frac{dy}{dx}$ in terms of t . 2
- (b) (i) Find the angle of projection of the stone. 2
- (ii) By considering $\frac{dy}{dx}$ find the value of t when the stone is moving at 45° below the horizontal. 2
11. A uniform lamina is bounded by the curve $y=x^3$, the line $x=4$ and the x -axis.



Find the coordinates of the centre of mass of the lamina.

4


[Turn over

12. An aircraft flies 1080 km due east from Glasgow to Copenhagen in a time of $2\frac{1}{4}$ hours.

The aircraft sets a course on a bearing of 100° and the speed of the aircraft in still air is 450 km h^{-1} .

- (a) Calculate the magnitude and direction of the wind. 3
- (b) (i) Given that the velocity of the wind remains constant, explain why the return journey will take longer. 1
- (ii) Calculate how much longer the return journey will take, giving your answer to the nearest minute. 4


13. A glass bowl is modelled by rotating the curve $y = e^{\frac{x}{12}}$ between $x = 15$ and $x = 30$ through 2π radians about the x -axis as shown in the diagram.



- (a) Find the volume of the bowl. 3
- (b) A line is to be put on the bowl to indicate when it is half full.
How far above the base of the bowl should this line be marked? 3

14. A block of weight W is placed on a rough inclined plane at an angle θ to the plane.

It can be held on the point of slipping down the plane by a force P acting parallel to the plane or a horizontal force Q as shown by the diagrams.



Prove that $P = \frac{QW}{Q \sin \theta + W \cos \theta}$.

7

15. A mass of 0.25 kg is attached to a horizontal spring of natural length 1 metre and modulus of elasticity 20 newtons. The spring is stretched and then released. It experiences a resistive force of magnitude $6v$ newtons, where v is the velocity of the mass.

- (a) Show that the subsequent motion satisfies the second order differential equation

$$\frac{d^2x}{dt^2} + 24 \frac{dx}{dt} + 80x = 0.$$

2

- (b) Solve this second order differential equation given that the mass is released from rest with an extension in the spring of 0.2 m.

6

- (c) Show that the acceleration is equal to zero when $t = \frac{1}{16} \ln 5$ seconds and find the displacement at this time.

3

[Turn over for next question]

16. A ball is projected from an origin on horizontal ground with speed $V \text{ m s}^{-1}$ at an angle of elevation of θ and moves freely under gravity. It passes through a point which is x metres horizontally from the origin at a height y metres above the ground.

(a) Show that the trajectory of the particle has equation

$$y = x \tan \theta - \frac{gx^2}{2V^2} (1 + \tan^2 \theta).$$

(Note that $\sec^2 \theta = 1 + \tan^2 \theta$)

3

- (b) The ball is at a vertical height of h metres when it has travelled $4h$ metres horizontally.


It is again at a height of h metres when it has travelled a further h metres horizontally.

Determine the angle of projection θ .

5

17. A light inextensible string of length r metres has one end attached to a fixed point O and the other end is attached to a particle of mass m kilograms.

From its equilibrium position, the particle is given a horizontal velocity $u \text{ m s}^{-1}$, as shown in the diagram.



- (a) (i) Show that the tension, T , in the string can be expressed as

$$T = \frac{mu^2}{r} + mg(3 \cos \theta - 2)$$

where θ is the angle between the string and the downward vertical through O.

4

- (ii) Determine a condition for u in terms of r and g , so that the particle executes a complete circle.

2

- (b) Given that the value of u is $2\sqrt{rg}$, find an expression in terms of r for the height of the particle above its starting position when the string goes slack.

3

[END OF QUESTION PAPER]